Entangled cluster states of four interacting spins with frustrated quasiparticles

Tieyan Si
Academy of Fundamental and Interdisciplinary Sciences,
Harbin Institute of Technology, Harbin, 150080, China
(Dated: April 2, 2013)

We constructed a four spin-1/2 model on a triangular star lattice to study the quantum statistical behavior of the frustrated triple-spin clusters and double-spin clusters. The double-spin cluster behaves as boson in zero-energy state and the highest excited state. While in the ground state and other intermediate excited states, the double spin cluster shows a hybrid statistical factor of fermion, boson and exotic quasiparticles. We find a definition for computing the statistical factor between two triple-spin plaquette quasiparticles. The statistical matrix of exchanging two triple-spin clusters is far beyond fermion and boson. The eigenstates of this model are maximal entangled quantum states like Greenberger-Horne-Zeilinger state and Yeo-Chua's genuine four-qubit entangled state. The zero-energy state is a generalized singlet state of two double-spin clusters. While the highest excited state is generalized symmetric state of two double-spin clusters. The quantum entanglement between two double-spin clusters is relatively trivial in comparison with that of triple-spin cluster state. The von Neumann entropy of triple-spin cluster state is much higher. Majonana fermion has interesting physical behavior in strongly interacted fermion system. This spin model is not directly related to Majorana fermions. So we represent Majorana fermions by string operator of spins following the inverse Jordan-Wigner transformation. In this representation, the Majorana fermions do not behaves like independent free operators but entangled with one another. Exchanging two Majorana fermions not only changed the output wave function of the two fermions, but also rearranged all the rest components of the collective wave function. This property is showed by computing the statistical matrix of Majorana fermions for exchanging two particles. The output wave function of Majorana fermions is collectively entangled quantum state. The exotic statistical matrix maybe useful for constructing quantum gate. The current quantum information technology maybe can implement this four particle model.

PACS numbers: 05.30.Pr, 03.67.Lx, 73.43.Cd

arXiv:1205.1052v3 [quant-ph] 1 Apr 2013

I. INTRODUCTION

Entangled quantum states is the central source of quantum computation. Topological ordered states as a promising error-correcting quantum code for fault-tolerant quantum computation has aroused broad theoretical research interest [1–12]. There had been many proposals for experimentally studying quasiparticle in Kitaev toric code model [13][14] as well as Kitaev honeycomb [15][16]. The laser excited Rydberg atoms in optical lattice is an effective quantum simulator [17] of Kitaev toric code model and color code model [18]. The solid material Na_2IrO_3 is also possible to map Kitaev honeycomb model into reality [19].

Majorana fermion appears as a promising candidate for topological quantum computation. It can be constructed by ordinary electron and hole in p-wave superconductor or quantum wire [20][21]. Direct observation of Majorana fermion is likely put into practice in quantum wire [22], quantum dots [23], trapped ions [24] and vortex of atomic fermi gases [25].

On the other route, quantum optics had been very successful to manipulate a small number of qubits. Greenberger-Horne-Zeilinger state has been generated in laboratory [26], so did the W-state of four qubits [27]. Yeo-Chua's genuine four-qubit entangled state is also promising for experiment [28]. The current quantum information technology can implement these entangled eigenstates by cross-Kerr nonlinearity [29]. The exciting condensed matter systems above are somewhat too complicate for a quantum optical implementation. Thus we constructed a four-spin model with frustrated quasiparticles. It inherits the exotic physics of strongly correlated many body system, in the meantime it is convenient for quantum optical research.

The strongly coupled four spins with frustrated quasiparticle is built on a triangular star lattice. The coupling rule between neighboring spins follows Kitaev honeycomb model. However we add an anti-ferromagnetic coupling between neighboring plaquette operators which sits right at the center of plaquette. This introduced the geometric frustration of the three quasiparticles. A frustrated system has large degenerated Hilbert space and nontrivial topological order [30]. Geometric frustration is the intrinsic reason for the exotic statistics of quasiparticles in this model.

The article is organized as following: In section II, we solved the triangular star model by diagonalizing the Hamiltonian matrix. The eigenenergy values and spin configurations of eigenstates are derived. In section III, we computed the statistical factor of double-spin clusters and triple-spin clusters. In section IV, we analyzed the symmetry of zero energy state and the highest excited state and computed the von Neumann entropy of the triple-spin cluster state. In section V, we computed the permutation matrix of two Majorana fermion operators on ground states. Exchanging two Majorana fermions goes far beyond fermionic statistics. The section VI is summary, we introduced the straight generalization of this triangular star model to a frustrated Kitaev honeycomb model. The appendix includes more details.

II. THE ENTANGLED STATES OF TWO DOUBLE-SPIN CLUSTERS IN THE TRIANGULAR STAR MODEL

In the triangular star model, the four particles are placed on the vertices of a triangular star lattice(Fig. 1). The triangular star has three independent triangular plaquette. The four particles coupled to each other following Kitaev honeycomb model. We added an antiferromagnetic coupling between the nearest neighboring plaquette on the Hamiltonian,

$$\begin{aligned} H_a &= J_x\sigma_1^x\sigma_3^x + J_y\sigma_1^y\sigma_2^y + J_z\sigma_2^z\sigma_3^z + J_x\sigma_2^x\sigma_4^x + J_y\sigma_3^y\sigma_4^y \\ &+ J_z\sigma_1^z\sigma_4^z + J_p\hat{S}_{\mathbf{1}}\hat{S}_{\mathbf{2}} + J_p\hat{S}_{\mathbf{2}}\hat{S}_{\mathbf{3}} + J_p\hat{S}_{\mathbf{3}}\hat{S}_{\mathbf{1}}. \end{aligned} \tag{1}$$

The three plaquette operators commute with the Hamiltonian. They are conserved quantum operators,



(a) **(b)**

FIG. 1: (a) The triangular star model. Quasiparticle sits right at the center of each triangle. (b) The eigenenergy levels of the triangular star model with respect to the parameter setting of ($J_p = 2J_x,\ J_z = 2J_x,\ J_y = 2J_x$).

$$\hat{S}_{\mathbf{1}} = \sigma_1^z\sigma_2^x\sigma_3^y,\ \ \hat{S}_{\mathbf{2}} = \sigma_4^z\sigma_2^y\sigma_3^x,\ \ \hat{S}_{\mathbf{3}} = \sigma_1^x\sigma_2^z\sigma_4^y. \tag{2}$$

The three conserved plaquette operator divide the total Hilbert space into three sectors. Each plaquette operator has eigenvalues +1 and −1 within its Hilbert space, $\hat{S}_\alpha|\psi\rangle = \pm 1|\psi\rangle$. Every triangular plaquette operator defines an effective Ising spin within each sector. In the Kitaev honeycomb model, the ground state chooses a homogeneous gauge pattern. All plaquette operators take the same eigenvalue. As there is antiferromagnetic coupling between two plaquette, the ground state is no longer the homogeneous gauge pattern. The ground state bear the frustrated gauge pattern.

We first solve the model by diagonalizing the Hamiltonian matrix. The spin operators take a sixteen dimensional representation,

$$\begin{aligned} \sigma_1^\mu &= \hat{\sigma}_1^\mu \otimes \mathbf{I}_2 \otimes \mathbf{I}_3 \otimes \mathbf{I}_4, \quad \sigma_2^\mu = \mathbf{I}_1 \otimes \hat{\sigma}_2^\mu \otimes \mathbf{I}_3 \otimes \mathbf{I}_4, \\ \sigma_3^\mu &= \mathbf{I}_1 \otimes \mathbf{I}_2 \otimes \hat{\sigma}_3^\mu \otimes \mathbf{I}_4, \quad \sigma_4^\mu = \mathbf{I}_1 \otimes \mathbf{I}_2 \otimes \mathbf{I}_3 \otimes \hat{\sigma}_4^\mu, \end{aligned} \tag{3}$$

where $\hat{\sigma}_i^\mu$ are the conventional Pauli matrices and $\mathbf{I}_i$ is the 2×2 identity matrix. The symbol ⊗ denotes direct product. The eigenvalues of the sixteen dimensional Hamiltonian matrix lead to eight discrete energy levels,

$$\begin{aligned} E_p^{\pm} &= 3J_p \pm 2\sqrt{J_x^2 + J_y^2 + J_z^2}, \\ E_\mu^{\pm} &= -J_p \pm 2J_\mu, \qquad \mu = x, y, z. \end{aligned} \tag{4}$$

Each energy level has two fold degeneracy. The energy levels are listed in Fig. 1 (b). The eigenenergy and eigenstates are computed directly from the Hamiltonian matrix. The newly added coupling terms of two plaquette operators commute with Hamiltonian. It does not modify the physics of Kitaev honeycomb model. The vortex excitation in the triangular plaquette represents the same type of quasiparticle excitation of Kitaev honeycomb model. The main different character of this triangle star model from Kitaev honeycomb model is the neighboring quasiparticles are now antiferromagnetically coupled to each other. So the ground state is the minimally frustrated quasiparticle state.

The two levels with higher eigenenergy $E_p^{\pm}$ correspond to excited states for which the eigenvalues of the three plaquette operators takes the same value $S_{\mathbf{1}} = S_{\mathbf{2}} = S_{\mathbf{3}} = \pm 1$. The three pairs of energy level with lower energy $E_\mu^{\pm}$ correspond to the frustrated gauge pattern. Only one pair of quasiparticles is not frustrated for $E_p^{\pm}$. If $J_x = J_y = J_z$, the six energy levels of $E_\mu^{\pm}$ would be degenerated.

If the quasiparticle coupling interaction J_p becomes zero, the triangular star model reduces to a finite Kitaev honeycomb model on triangular star lattice. For a stronger quasiparticle coupling interaction than the spin coupling,

$$J_p = 2J_x, \quad J_z = 2J_x, \quad J_y = 2J_x, \tag{5}$$

there exists four degenerated states with zero energy. Without losing important physics, we focus on this special parameter setting of Eq. (5) in the following. By representing the eigenvectors of the 16 dimensional Hamiltonian matrix by the 16 four-spin basis, we derived the spin configurations corresponding to the four zero energy states,

$$\begin{aligned} |o\rangle_1 &= |\Uparrow\Uparrow\rangle + |\Downarrow\Downarrow\rangle + 2|\circlearrowright\circlearrowleft\rangle + 2|\circlearrowleft\circlearrowright\rangle \\ &- 5|\circlearrowleft\circlearrowleft\rangle - 5|\circlearrowright\circlearrowright\rangle, \\ |o\rangle_2 &= |\Uparrow\circlearrowright\rangle + |\circlearrowright\Uparrow\rangle + |\circlearrowleft\Downarrow\rangle + |\Downarrow\circlearrowleft\rangle, \\ |o\rangle_3 &= |\Uparrow\circlearrowleft\rangle + |\circlearrowright\Downarrow\rangle + |\circlearrowleft\Uparrow\rangle + |\Downarrow\circlearrowright\rangle, \\ |o\rangle_4 &= |\Uparrow\Downarrow\rangle + |\Downarrow\Uparrow\rangle + 2|\circlearrowleft\circlearrowleft\rangle + 2|\circlearrowright\circlearrowright\rangle \\ &- |\circlearrowright\circlearrowleft\rangle - |\circlearrowleft\circlearrowright\rangle. \end{aligned} \tag{6}$$

Here the symbols in the quantum wave function represent the double spin configurations,

$$|\Uparrow\rangle = |\uparrow\uparrow\rangle,\ |\Downarrow\rangle = |\downarrow\downarrow\rangle,\ |\circlearrowright\rangle = |\uparrow\downarrow\rangle,\ |\circlearrowleft\rangle = |\downarrow\uparrow\rangle. \tag{7}$$

The eigenenergy of the states above is zero, i.e., $H|o\rangle_\mu = 0, \ \ \mu = 1, 2, 3, 4$. The zero energy state pair of $|o\rangle_2$ and

$|o\rangle_3$ comes from the minimal frustrated gauge pattern, i.e, $E_x^+ = -J_p + 2J_x = 0$. The two zero energy states of $|o\rangle_1$ and $|o\rangle_4$ correspond to the fully frustrated states $E_p^- = 3J_p - 6J_x = 0$.

The ground state has four fold degeneracy. We denote them as $|g\rangle_\mu, \;\; \mu = 1, 2, 3, 4.$,

$$\begin{aligned}
|g\rangle_1 &= |\Downarrow\circlearrowleft\rangle - |\Uparrow\circlearrowright\rangle + |\Uparrow\circlearrowleft\rangle - |\Downarrow\circlearrowright\rangle, \\
|g\rangle_2 &= |\Downarrow\Uparrow\rangle - |\Uparrow\Downarrow\rangle, \\
|g\rangle_3 &= |\circlearrowleft\Uparrow\rangle - |\circlearrowright\Downarrow\rangle + |\circlearrowleft\Downarrow\rangle - |\circlearrowright\Uparrow\rangle, \\
|g\rangle_4 &= |\circlearrowleft\circlearrowleft\rangle - |\circlearrowright\circlearrowright\rangle.
\end{aligned} \tag{8}$$

The eigenenergy of the four ground states are $E_g = -6J_x$, i.e., $H|g\rangle_\mu = -6J_x|g\rangle_\mu, \quad \mu = 1, 2, 3, 4$. The ground states are the minimally frustrated quasiparticle states. Two of them comes from $E_y^- = -J_p - 2J_y = -6J_x$. The other two states is for $E_z^- = -J_p - 2J_z = -6J_x$. The spin configuration of the two states $|g\rangle_1$ and $|g\rangle_3$ is the superposition of three-up-one-down state and three-down-one-up state. The two states of $|g\rangle_2$ and $|g\rangle_4$ is the superposition state of four spin basis with two-up spins and two-down spins. If we flip the spins of all of the four ground states, it generates a minus sign upon the original wave function. So the ground state breaks Z_2 symmetry.

The whole Hilbert space of eigenstates can be classified into two classes. The first class breaks the Z_2 symmetry. This class only includes the ground state and the four degenerated states with respect to $2J_x$. The other class keeps Z_2 symmetry. This class includes the zero energy states, the highest excited states and all the rest excited states.

The energy level $\omega = 12J_x$ is the highest energy level corresponding to the fully frustrated quasiparticle state. The spin configuration corresponding to this two-fold degenerated level is $|e\rangle_{15}$ and $|e\rangle_{16}$,

$$\begin{aligned}
|e\rangle_{15} &= |\Uparrow\Uparrow\rangle + |\Downarrow\Downarrow\rangle + 2|\circlearrowright\circlearrowleft\rangle + 2|\circlearrowleft\circlearrowright\rangle \\
&+ |\circlearrowleft\circlearrowleft\rangle + |\circlearrowright\circlearrowright\rangle, \\
|e\rangle_{16} &= |\Uparrow\Downarrow\rangle + |\Downarrow\Uparrow\rangle + 2|\circlearrowright\circlearrowright\rangle + 2|\circlearrowleft\circlearrowleft\rangle \\
&+ 5|\circlearrowleft\circlearrowright\rangle + 5|\circlearrowright\circlearrowleft\rangle,
\end{aligned} \tag{9}$$

The eigenstates of this model can be viewed as entangled quantum states of two double-spin clusters. For example, the nearest excited state above the zero energy state has four-fold degeneracy. The spin configuration of the four states are

$$\begin{aligned}
|e\rangle_{11} &= -|\Uparrow\Uparrow\rangle + |\Downarrow\Downarrow\rangle, \\
|e\rangle_{12} &= -|\Uparrow\circlearrowright\rangle - |\Uparrow\circlearrowleft\rangle + |\Downarrow\circlearrowright\rangle + |\Downarrow\circlearrowleft\rangle, \\
|e\rangle_{13} &= -|\circlearrowright\Uparrow\rangle + |\circlearrowright\Downarrow\rangle - |\circlearrowleft\Uparrow\rangle + |\circlearrowleft\Downarrow\rangle, \\
|e\rangle_{14} &= -|\circlearrowright\circlearrowleft\rangle + |\circlearrowleft\circlearrowright\rangle.
\end{aligned} \tag{10}$$

The corresponding eigenenergy with respect to $\{|e\rangle_{11}$, $|e\rangle_{12}$, $|e\rangle_{13}$, $|e\rangle_{14}\}$ is $E = 2J_x$. The spin configurations of $|e\rangle_{12}$ and $|e\rangle_{13}$ is the superposition of three-up-one-down and three-down-one-up. The state $|e\rangle_{11}$ can be viewed as a dual state of the well-known Greenberger-Horne-Zeilinger state for four qubit[26],

$$|GHZ\rangle = |\Uparrow\Uparrow\rangle + |\Downarrow\Downarrow\rangle. \tag{11}$$

$|e\rangle_{14}$ is a singlet pair state of two double-spin clusters . The other two states, $|e\rangle_{12}$ and $|e\rangle_{13}$, can be viewed as generalized W-state of four qubits [27],

$$|W\rangle = (|\Uparrow\circlearrowright\rangle + |\Uparrow\circlearrowleft\rangle + |\circlearrowright\Uparrow\rangle + |\circlearrowleft\Uparrow\rangle). \tag{12}$$

The two degenerated states of energy level $E = -4J_x$ are $|e\rangle_9\}$ and $\{|e\rangle_{10}$, they both bear simmilar state structure as W-state,

$$\begin{aligned}
|e\rangle_9 &= |\Uparrow\circlearrowright\rangle - |\circlearrowright\Uparrow\rangle - |\circlearrowleft\Downarrow\rangle + |\Downarrow\circlearrowleft\rangle, \\
|e\rangle_{10} &= |\Uparrow\circlearrowleft\rangle - |\circlearrowleft\Uparrow\rangle - |\circlearrowright\Downarrow\rangle + |\Downarrow\circlearrowright\rangle.
\end{aligned} \tag{13}$$

These eigenstates are genuine entangled states of four spins. The zero energy states and highest excited states, $\{|o\rangle_1, |o\rangle_4\}$ and $\{|e\rangle_{15}, |e\rangle_{16}\}$, can be classified into the same class as Yeo-Chua's genuine four-qubit entangled state [28],

$$\begin{aligned}
|\chi^{00}\rangle &= |\Downarrow\Downarrow\rangle - |\Downarrow\Uparrow\rangle - |\circlearrowleft\circlearrowleft\rangle + |\circlearrowleft\circlearrowright\rangle \\
&+ |\circlearrowright\circlearrowleft\rangle + |\circlearrowright\circlearrowright\rangle + |\Uparrow\Downarrow\rangle + |\Uparrow\Uparrow\rangle.
\end{aligned} \tag{14}$$

The eigenstates of this model can be implemented by the same operation as that for generating Yeo-Chua's genuine four-qubit entangled state using cross-Kerr nonlinearity [29]. These states can be mapped into graph states following the strategy of Ref. [31], the observable operators in graph state theory has a physical meaning in this quantum model. Usually Yeo-Chua's genuine four-qubit entangled state or W-state is constructed without a Hamiltonian, but here we can derive the eigenenergy of these states. One can decompose an arbitrary entangled state as the superposition of these eigenstates. The weight of each eigenstate is marked by its eigenvalue. This offers us a new angle to see the internal structure of the quantum entanglement.

The Wootters's concurrence provide a convenient way to quantify quantum entanglement of four-qubit states[32],

$$\tau_{abcd} = |\langle\psi|\hat{\sigma}_a^y \otimes \hat{\sigma}_b^y \otimes \hat{\sigma}_c^y \otimes \hat{\sigma}_d^y|\psi^*\rangle|^2. \tag{15}$$

This concurrence operator is happened to be a conserved quantum operator of this triangular star model. The product of any two plaquette operators is a string of four identical spin operators,

$$\begin{aligned}
\hat{S}_{\mathbf{1}}\hat{S}_{\mathbf{2}} &= \sigma_1^z\sigma_2^z\sigma_3^z\sigma_4^z, \quad \hat{S}_{\mathbf{2}}\hat{S}_{\mathbf{3}} = \sigma_1^x\sigma_2^x\sigma_4^x\sigma_3^x, \\
\hat{S}_{\mathbf{3}}\hat{S}_{\mathbf{1}} &= \sigma_1^y\sigma_2^y\sigma_3^y\sigma_4^y.
\end{aligned} \tag{16}$$

The plaquette operators keep an arbitrary ground state vector within ground state. For example, the operation of plaquette operator $\hat{S}_{\mathbf{1}}$ on the vector ground state, $\Psi_{[g_{1234}]} = [\ |g\rangle_1,\ |g\rangle_2\ , |g\rangle_3\ , |g\rangle_4\ ]^T$, gives a matrix,

$$\hat{S}_{\mathbf{1}}\Psi_{[g_{1234}]} = \begin{bmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & -i \\ -i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{bmatrix} \begin{bmatrix} |g\rangle_1 \\ |g\rangle_2 \\ |g\rangle_3 \\ |g\rangle_4 \end{bmatrix}. \tag{17}$$

The operation of concurrence operator reads, $\hat{\tau} = \hat{S}_{\mathbf{3}}\hat{S}_{\mathbf{1}}$,

$$\hat{\tau}\Psi_{[g_{1234}]} = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} |g\rangle_1 \\ |g\rangle_2 \\ |g\rangle_3 \\ |g\rangle_4 \end{bmatrix}. \quad (18)$$

III. THE STATISTICAL PROPERTY OF DOUBLE-SPIN CLUSTER STATES AND PLAQUETTE QUASIPARTICLES

As all know, switching two fermions would generate a negative sign in front of the collective wave function. What would happen if it switches two double-spin clusters? Usually a particle composed of two fermions behaves like a boson. However some of the eigenstates of the triangular star model behaves like a fermionic wave function of two double-spin clusters. We define a permutation operator, $P[12;34]$,

$$P[12;34]\Psi(1234) = \Psi(3412) = \eta\Psi(1234). \quad (19)$$

If $\eta = +1$, the cluster state is bosonic state. If $\eta = -1$, the cluster state is fermionic state. If η bear more complex structure, we might call the clusters as exotic quasiparticles.

The four degenerated ground states include one femionic cluster state and one bosonic cluster states,

$$P[12;34]|g\rangle_2 = (-1)|g\rangle_2, \;\; P[12;34]|g\rangle_4 = (+1)|g\rangle_4. (20)$$

The rest two ground states can map into each other by the permutation operator,

$$P[12;34]|g\rangle_1 = |g\rangle_3, \qquad P[12;34]|g\rangle_3 = |g\rangle_1. \quad (21)$$

So we define a four dimensional vector of ground cluster state,

$$\Psi_{[g_{(1234)}]} = [\ |g\rangle_1,\ |g\rangle_2\ , |g\rangle_3\ , |g\rangle_4\ ]^T, \quad (22)$$

then the permutation operates on the vector states,

$$P[12;34]\Psi_{[g_{(1234)}]} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} |g\rangle_1 \\ |g\rangle_2 \\ |g\rangle_3 \\ |g\rangle_4 \end{bmatrix}. \quad (23)$$

If we select two states out of the four ground states as basis for quantum computation, the statistical factor of $\Psi_{[g_1,g_3]} = [\ |g\rangle_1,\ |g\rangle_3\ ]^T$ is a Pauli matrix $\eta = \sigma^x$. For the vector wave function of $\Psi_{[g_2,g_4]} = [\ |g\rangle_2,\ |g\rangle_4\ ]^T$, the statistical factor of $\Psi_{[g_2,g_4]}$ is another Pauli matrix $\eta = \sigma^z$. The statistical property of the nearest excited state above zero energy is similar to ground state,

$$\begin{aligned} P[12;34]|e\rangle_{11} &= |e\rangle_{11}, \quad P[12;34]|e\rangle_{14} = -|e\rangle_{14}. \\ P[12;34]|e\rangle_{12} &= |e\rangle_{13}, \quad P[12;34]|e\rangle_{13} = |e\rangle_{12}. \end{aligned} \quad (24)$$

The four degenerate zero energy states and the highest excited states are all bosonic cluster states,

$$\begin{aligned} P[12;34]|o\rangle_i &= |o\rangle_i, \;\; i = 1,2,3,4; \\ P[12;34]|e\rangle_j &= |e\rangle_j, \;\; j = 15,16. \end{aligned} \quad (25)$$

The first excited state has two degenerated states. Permutating two clusters produce a statistical factor beyond boson and fermions,

$$\begin{aligned} P[12;34]|e\rangle_9 &= e^{i\pi}[\sigma_3^x\sigma_4^x]|e\rangle_{10}, \\ P[12;34]|e\rangle_{10} &= e^{i\pi}[\sigma_3^x\sigma_4^x]|e\rangle_9, \end{aligned} \quad (26)$$

For the vector wave function of the first excited states, $\Psi_{[e_9,e_{10}]} = [\ |e\rangle_9,\ |e\rangle_{10}\ ]^T$, the statistical factor of double spin clusters is

$$\eta^e = e^{i\pi}\sigma_3^x\sigma_4^x \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}. \quad (27)$$

The three plaquette operator define three quasiparticles. These quasiparticles are eigen-excitations of the quantum model. To find out if the quasiparticles are fermion or boson, we define the same permutation operator as above,

$$P[S_{\mathbf{1}};S_{\mathbf{2}}] \;=\; P[1;4]\Psi(1234) = \eta_{[S_{\mathbf{1}};S_{\mathbf{2}}]}\Psi(1234). \quad (28)$$

The output of the permutation operator $P[S_{\mathbf{1}};S_{\mathbf{2}}]$ on the four dimensional vector ground states $\Psi_{[g(1234)]}$ is a nontrivial statistical matrix,

$$\eta^g_{[S_{\mathbf{1}};S_{\mathbf{2}}]} = \begin{bmatrix} 0 & -\sigma_3^x & 0 & \sigma_1^x \\ 0 & 0 & 0 & -1 \\ 0 & \sigma_4^x & 0 & -\sigma_3^x \\ 0 & -1 & 0 & 0 \end{bmatrix}. \quad (29)$$

Following the same procedure, one may find the statistical matrix of $P[S_{\mathbf{2}};S_{\mathbf{3}}] = P[1;3]\Psi(1234)$, $P[S_{\mathbf{1}};S_{\mathbf{3}}]$, and so on. If quasiparticle $S_{\mathbf{1}}$ complete a closed trajectory and returns to its starting point, then the statistical matrices satisfies the relationship,

$$P[S_{\mathbf{3}};S_{\mathbf{2}}]P[S_{\mathbf{2}};S_{\mathbf{1}}]P[S_{\mathbf{1}};S_{\mathbf{3}}]P[S_{\mathbf{1}};S_{\mathbf{2}}] = I, \quad (30)$$

where I is a four dimensional identity matrix.

In the first excited state, $\Psi_{[e_9,e_{10}]} = [\ |e\rangle_9,\ |e\rangle_{10}\ ]^T$, the triple-spin cluster is a hybrid of fermion and boson. The fermionic part and bosonic part of the first state can be separated as

$$\begin{aligned} A_9 &= |\Uparrow\circlearrowright\rangle + |\Downarrow\circlearrowleft\rangle, \qquad B_9 = -|\circlearrowright\Uparrow\rangle - |\circlearrowleft\Downarrow\rangle, \\ A_{10} &= -|\circlearrowleft\Uparrow\rangle - |\circlearrowright\Downarrow\rangle, \;\; B_{10} = |\Uparrow\circlearrowleft\rangle + |\Downarrow\circlearrowright\rangle. \end{aligned} \quad (31)$$

The operation of permutation operator $P[S_{\mathbf{1}};S_{\mathbf{2}}]$ on the first excited states, $|e\rangle_9 = A_9 + B_9$, $\;|e\rangle_{10} = A_{10} + B_{10}$, gives us,

$$\begin{aligned} P[S_{\mathbf{1}};S_{\mathbf{2}}]|e\rangle_9 &= -A_{10} + B_9, \\ P[S_{\mathbf{1}};S_{\mathbf{2}}]|e\rangle_{10} &= -A_9 + B_{10}, \end{aligned} \quad (32)$$



FIG. 2: The permutating steps for plaquette quasiparticle $S_{\mathbf{1}}$ to complete a circle around the center.

We define the statistical factor of the vector wave function of $\Psi_{[e_9,e_{10}]} = [\ |e\rangle_9,\ |e\rangle_{10}\ ]^T$ as

$$P[S_{\mathbf{1}};S_{\mathbf{2}}]\Psi_{[e_9,e_{10}]} = \eta^e_{[S_{\mathbf{1}};S_{\mathbf{2}}]}\Psi_{[e_9,e_{10}]}. \tag{33}$$

The statistical factor can be derived by matrix algebra,

$$\eta^e_{[S_{\mathbf{1}};S_{\mathbf{2}}]} = -\left[\begin{array}{cc} \sigma^x_2\sigma^x_4[I-\sigma^x_2\sigma^x_4]^{-1} & [I-\sigma^x_1\sigma^x_3]^{-1} \\ [I-\sigma^x_2\sigma^x_4]^{-1} & \sigma^x_1\sigma^x_3[I-\sigma^x_1\sigma^x_3]^{-1} \end{array}\right] \tag{34}$$

here I is 2 by 2 identical matrix. So the plaquette quasiparticle has complicate statistical behavior. This exotic property may be used to construct non-trivial quantum logical gate.

IV. THE TRANSFORMATION BETWEEN SYMMETRIC CLUSTER STATE AND ANTISYMMETRIC CLUSTER STATE

The conventional spin 1/2 particles may form a antisymmetric state $(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$ and a symmetric state $(|\uparrow\uparrow\rangle + |\downarrow\downarrow\rangle)$. Usually the singlet state leads to lower energy. The triplet leads to higher energy. A singlet can be transformed into a triplet state by flipping the second spin or the first spin,

$$[\ \sigma^x_2|\uparrow\downarrow\rangle - e^{i\pi}\sigma^x_2|\downarrow\uparrow\rangle\ ] \quad => \quad [\ |\uparrow\uparrow\rangle + |\downarrow\downarrow\rangle\ ]. \tag{35}$$

The singlet state and triplet state above is defined for two spins. While the spin configuration of the triangle star model includes four spins. A similar transformation rule as that for a conventional singlet state and a triplet state also exist between the zero energy state and the highest energy states of this triangular star model. Here is we need to flip a pair of spins. Based on the spin configuration of zero energy states and the highest excited states, one may extract two unit configuration out of the complete states,

$$\begin{aligned} A &= |\Uparrow\Uparrow\rangle + 2|\circlearrowright\circlearrowleft\rangle + 2|\circlearrowleft\circlearrowright\rangle + |\Downarrow\Downarrow\rangle, \\ B &= 5|\circlearrowleft\circlearrowleft\rangle + 5|\circlearrowright\circlearrowright\rangle. \end{aligned} \tag{36}$$

Both the zero energy state and the highest excited state are linear combination of the two units,

$$|o\rangle_1 = A - B, \quad |e\rangle_{16} = [\sigma^x_3\sigma^x_4]A - e^{i\pi}[\sigma^x_3\sigma^x_4]B. \tag{37}$$

The same algebra relation also exist between the state $|o\rangle_4$ and $|e\rangle_{15}$. The two unit spin configurations of $|o\rangle_4$ and $|e\rangle_{15}$ are different from that of $|o\rangle_1$ and $|e\rangle_{16}$.

Both $|o\rangle_1$ and $|e\rangle_{16}$ are bosonic state of double spin clusters. However the statistical factor of triple-spin cluster state is beyond fermion or boson statistics. Any two of the three plaquette always share two spins, so one can not cut one triple-spin cluster out of the four spins to get an independent three-spin wave function. We use the reduced density operator to represent the triple-spin cluster state. The density operator for the zero energy state $|o\rangle_1$ of quasiparticle $\hat{S}_{\mathbf{1}} = \sigma^z_1\sigma^x_2\sigma^y_3$ reads $\rho_{S_{\mathbf{1}}}(o_1) = Tr_{[4]}\left[[|o\rangle_{11}\langle o|]_{[1234]}\right]$. A detail calculation yields

$$\rho_{S_{\mathbf{1}}}(o_1) = \begin{pmatrix} 1 & 0 & 0 & 2 & 0 & -5 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 25 & 0 & -10 & 0 & 0 & -5 \\ 2 & 0 & 0 & 4 & 0 & -10 & 0 & 0 \\ 0 & 0 & -10 & 0 & 4 & 0 & 0 & 2 \\ -5 & 0 & 0 & -10 & 0 & 25 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -5 & 0 & 2 & 0 & 0 & 1 \end{pmatrix}. \tag{38}$$

We also derived the density operator of $S_{\mathbf{1}}$ in the highest excited state $|e\rangle_{16}$,

$$\rho_{S_{\mathbf{1}}}(e_{16}) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 & 5 & 0 & 0 & 0 \\ 0 & 2 & 4 & 0 & 10 & 0 & 0 & 0 \\ 0 & 0 & 0 & 25 & 0 & 10 & 5 & 0 \\ 0 & 5 & 10 & 0 & 25 & 0 & 0 & 0 \\ 0 & 0 & 0 & 10 & 0 & 4 & 2 & 0 \\ 0 & 0 & 0 & 5 & 0 & 2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \tag{39}$$

The two density operators of quasiparticle $S_{\mathbf{1}}$ have the same eigenvalues, $\vec{\lambda} = [1/2, 1/2, 0, 0, 0, 0, 0, 0]$. Only the two non-zero eigenvalues, $\lambda_1 = 1/2$ and $\lambda_2 = 1$, contribute to von Neumann entropy,

$$S(\rho) = -\lambda_1 Log[\lambda_1] - \lambda_2 Log[\lambda_2] = 1. \tag{40}$$

von Neumann entropy quantifies the quantum entanglement of the triple-spin quasiparticle. The von Neumann entropy of quasiparticle $S_{\mathbf{2}} = \sigma^y_2\sigma^x_3\sigma^z_4$ and $S_{\mathbf{3}} = \sigma^y_4\sigma^x_1\sigma^z_2$ are the same as $S_{\mathbf{1}}$. The density operator of $S_{\mathbf{2}}$ for the zero energy state $|o\rangle_1$ reads

$$\rho_{S_{\mathbf{2}}}(o_1) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & -5 & 2 & 0 \\ 0 & 4 & -10 & 0 & 0 & 0 & 0 & 2 \\ 0 & -10 & 25 & 0 & 0 & 0 & 0 & -5 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -5 & 0 & 0 & 0 & 0 & 25 & -10 & 0 \\ 2 & 0 & 0 & 0 & 0 & -10 & 4 & 0 \\ 0 & 2 & -5 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \tag{41}$$

Correspondingly the density operator for the highest excited states $|e\rangle_{16}$ is

$$\rho_{S_{\mathbf{2}}}(e_{16}) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 25 & 10 & 0 & 5 & 0 & 0 & 0 \\ 0 & 10 & 4 & 0 & 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 2 & 5 & 0 \\ 0 & 5 & 2 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 4 & 10 & 0 \\ 0 & 0 & 0 & 5 & 0 & 10 & 25 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \quad (42)$$

The reduced density operator of $S_{\mathbf{3}}$ is originally based on the spatial order [124]. If we put the last spin No. [4] in front of [12], the modified spatial order of the three spins [412] generates the same density operator as that of $S_{\mathbf{2}}$. The eigenvalues of $\rho_{S_{\mathbf{3}[124]}}(o_1)$ is the same as that $\rho_{S_{\mathbf{3}[412]}}(o_1)$. Thus the density operator of $S_{\mathbf{3}}$ is equivalent to $S_{\mathbf{2}}$, i.e., $\rho_{S_{\mathbf{3}}}(o_1) = \rho_{S_{\mathbf{2}}}(o_1)$, $\rho_{S_{\mathbf{3}}}(e_{16}) = \rho_{S_{\mathbf{2}}}(e_{16})$.

Both the eigenvalues of $S_{\mathbf{2}}$ and $S_{\mathbf{3}}$ is the same as the quasiparticle $S_{\mathbf{1}}$. The von-Neumann entropy of the density operator of the three quasiparticles are the same as well. Since the three quasiparticles intersect with one another, it is not a surprise if they have the same degree of quantum entanglement.

V. THE STATISTICAL FACTOR OF MAJORANA FERMION OPERATORS ACTING ON GROUND STATE

Kitaev honeycomb model was solved by two different approaches, the first one is representing a spin operator by two Majorana fermions [1]. The second approach is using Jordan-Wigner transformation [4]. Jordan-Wigner transformation maps spin to fermion or vice versa [33]. Here we combine the two different approaches [7] to represent Majorana fermion by spins. These Majorana fermion operators are composite operators, it generates complicate entangled states.

The explicit formulation of inverse Jordan-Wigner transformation depends on the spatial order of the four particles. We squeeze the triangular star lattice to a one dimensional chain by keeping the topology of interacting bonds invariant. The indices $('1','2','3','4')$ in the Hamiltonian Eq. (1) are the name of the four particles instead of its spatial ordering in the one dimensional chain. We denote the four spatial positions along the one dimensional chain as $(\mathbf{i}_1 < \mathbf{i}_2 < \mathbf{i}_3 < \mathbf{i}_4)$(Fig. 3). If we place particle $'1'$ on $\mathbf{i}_4$, the other three particles would sit on the rest three sites $(\mathbf{i}_1, \mathbf{i}_2, \mathbf{i}_3)$ before $\mathbf{i}_4$. For a special case of the spatial ordering of the four particles [1423], i.e., $(\mathbf{i}_1 = 1,\ \mathbf{i}_2 = 4,\ \mathbf{i}_3 = 2,\ \mathbf{i}_4 = 3)$, the inverse Jordan-Wigner transformation defines the spin representation of eight Majorana fermions,

$$\begin{aligned} \psi_1 &= \sigma_1^y, \qquad \psi_4 = \sigma_4^x \sigma_1^z, \qquad \psi_2 = \sigma_2^y \sigma_1^z \sigma_4^z, \\ b_1 &= -\sigma_1^x, \quad\ b_4 = -\sigma_4^y \sigma_1^z, \quad\ b_2 = -\sigma_2^x \sigma_1^z \sigma_4^z, \\ \psi_3 &= \sigma_3^x \sigma_1^z \sigma_4^z \sigma_2^z, \quad\ b_3 = -\sigma_3^y \sigma_1^z \sigma_4^z \sigma_2^z. \end{aligned} \quad (43)$$



FIG. 3: The one dimensional scheme of the triangular star model. The initial spatial ordering of the four particle is [1423]. Then we braid the two particle at the second site and the third site, it leads to [1243]. The coupling bonds connecting the four particles remain the same as the original triangular star model.

The string operator of spins for Majorana fermion exactly fulfills the commutator of the original Majorana fermions, $b_i^\dagger = b_i$, $\psi_i^\dagger = \psi_i$, $\{b_i, b_j\} = \delta_{ij}$, $\{\psi_i, \psi_j\} = \delta_{ij}$, $\{b_i, \psi_j\} = 0$.

First we need to label the spatial position of the Majorana fermions. The inverse Jordan-Wigner transformation provides a natural way for labeling each Majorana fermion. The first Majorana fermion is represented by one spin operator, the second is expressed by the product of two spin operators, and so forth. If we exchange the position of two Majorana fermions, its representation would change spontaneously, so does its action on quantum state.

Then we introduce a vector of eight identical $|g\rangle_4$ states,

$$\overrightarrow{|g\rangle}_4 = [|g\rangle_4, |g\rangle_4, |g\rangle_4, |g\rangle_4, |g\rangle_4, |g\rangle_4, |g\rangle_4, |g\rangle_4]^T. (44)$$

We define a spatially ordered matrix operator of the eight Majorana fermions

$$\hat{F}_{[\mathbf{i}_1\mathbf{i}_2\mathbf{i}_3\mathbf{i}_4]} = diag[\psi_1, \psi_2, \psi_3, \psi_4, b_1, b_2, b_3, b_4]. \quad (45)$$

The subscribe number '1' of the fermion ψ_1 is the name of particle. ψ_1 at different position is represented by a string operator with different length. The subscribes of $\hat{F}_{[\mathbf{i}_1\mathbf{i}_2\mathbf{i}_3\mathbf{i}_4]}$ defines the spatial order of the four particle. For example, $\hat{F}_{[4231]}$ means particle No. $'1'$ sits in the fourth site $\mathbf{i}_4$. Then ψ_1 is represented by the product of four spin operators. This diagonal matrix operator always has the form $diag[\psi_1, \psi_2, \psi_3, \psi_4, b_1, b_2, b_3, b_4]$. The position of a Majorana fermions is represented by the length of spin representation. When we exchange two Majorana fermions, the string representation of the two Majorana fermions would exchange.

We operate the Majorana fermions on the ground state $|g\rangle_4$ to study their statistical matrix. The output vector of matrix operator $\hat{F}_{[\mathbf{i}_1\mathbf{i}_2\mathbf{i}_3\mathbf{i}_4]}$ on $\overrightarrow{|g\rangle}_4$ can be denoted as

$$\begin{aligned} \overrightarrow{|\psi_{\overleftarrow{g_4}}\rangle} \;=\; & [|\psi_{\overleftarrow{g_4}}\rangle_1, |\psi_{\overleftarrow{g_4}}\rangle_2, |\psi_{\overleftarrow{g_4}}\rangle_3, |\psi_{\overleftarrow{g_4}}\rangle_4, \\ & |\psi_{\overleftarrow{g_4}}\rangle_5, |\psi_{\overleftarrow{g_4}}\rangle_6, |\psi_{\overleftarrow{g_4}}\rangle_7, |\psi_{\overleftarrow{g_4}}\rangle_8,]^T. \end{aligned} \tag{46}$$

The action of the matrix operator is equivalent to a statistical matrix,

$$\hat{F}_{[\mathbf{i}_1\mathbf{i}_2\mathbf{i}_3\mathbf{i}_4]}\overrightarrow{|g\rangle}_4 \;=\; F_{[\mathbf{i}_1\mathbf{i}_2\mathbf{i}_3\mathbf{i}_4]}\overrightarrow{|\psi_{\overleftarrow{g_4}}\rangle}. \tag{47}$$

The output matrix of $F_{[\mathbf{i}_1\mathbf{i}_2\mathbf{i}_3\mathbf{i}_4]}$ describes the statistical behavior of the Majorana fermions. If it is simply $F = diag[1,1,1,1,1,1,1,1]$ after we exchange two Majorana fermions, that means the two Majorana fermions behaves like boson. If it is $F = diag[-1,-1,-1,-1,-1,-1,-1,-1]$, that means the two Majorana fermions behaves like ordinary fermions. If $F_{[\mathbf{i}_1\mathbf{i}_2\mathbf{i}_3\mathbf{i}_4]}$ is complicate matrix beyond the two above, the two Majorana fermions have exotic statistics.

We set the initial position of the eight Majorna fermions as $\hat{F}_{[1423]}$. Eq. (43) summarizes the spin representation of Majorana fermions for the spatial ordering of [1423]. We operate these string operators on $|g\rangle_4$ and express the output spin configurations by linear combination of other eigenstates,

$$\begin{aligned} |\psi_{\overleftarrow{g_4}}\rangle_1 &= \frac{-i}{2}(|o\rangle_3 + |e\rangle_{10}), \quad |\psi_{\overleftarrow{g_4}}\rangle_2 = \frac{-i}{2}(|o\rangle_2| + |e\rangle_9), \\ |\psi_{\overleftarrow{g_4}}\rangle_3 &= \frac{-1}{2}(|o\rangle_3 - |e\rangle_{10}), \quad |\psi_{\overleftarrow{g_4}}\rangle_4 = \frac{-1}{2}(|o\rangle_2 - |e\rangle_9), \\ |\psi_{\overleftarrow{g_4}}\rangle_5 &= \frac{1}{2}(|e\rangle_{12} - |g\rangle_1), \quad |\psi_{\overleftarrow{g_4}}\rangle_6 = \frac{1}{2}(|g\rangle_1 + |e\rangle_{12}), \\ |\psi_{\overleftarrow{g_4}}\rangle_7 &= i\frac{1}{2}(|g\rangle_3 - |e\rangle_{13}), \quad |\psi_{\overleftarrow{g_4}}\rangle_8 = i\frac{1}{2}(|e\rangle_{13} + |g\rangle_3). \end{aligned} \tag{48}$$

The eight dimensional output vector $\overrightarrow{|\psi_{\overleftarrow{g_4}}\rangle}$ is the initial output for the spatial ordering [1423]. The initial matrix for $\hat{F}_{[1423]}$ is an identity $F_{[1423]} = diag[1,1,1,1,1,1,1,1]$ since the exchanging operation is still not performed yet.

We take two different paths of permutation operation for the four particles. For the first path, we first exchange the second particle with the third particle, then exchange the third particle with the fourth particle. Here we introduce a permutation operator $B_{\mathbf{i}_2\mathbf{i}_3}$ for exchanging the particle at site $\mathbf{i}_2$ with that at $\mathbf{i}_3$, $B_{\mathbf{i}_2\mathbf{i}_3}[1423] = [1243]$,

$$F_{[1243]} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -i & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & i & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -i & 0 & 0 & 0 & 0 \end{pmatrix}. \tag{49}$$

$F_{[1243]}$ suggest that exchanging two Majorana femions does not only add a minus sign in front of the initial wave function. The we continue to perform the second permutation upon the first permutation, $B_{\mathbf{i}_3\mathbf{i}_4}[1243] = [1234]$. The final permutation matrix for the final ordering [1234] is

$$F_{[1234]} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -i & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & i & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & i & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -i & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & i & 0 & 0 \end{pmatrix}. \tag{50}$$

For the second path, we first exchange $[\mathbf{i}_3\mathbf{i}_4]$ and then exchange $[\mathbf{i}_2\mathbf{i}_3]$(Appendix E),

$$B_{\mathbf{i}_3\mathbf{i}_4}[1423] = [1432], \quad B_{\mathbf{i}_2\mathbf{i}_3}[1432] = [1342]. \tag{51}$$

The final permutation matrix for [1342] reads

$$F_{[1342]} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -i & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -i & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -i & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -i \end{pmatrix}. \tag{52}$$

Usually if we exchange Majorana fermions twice, the statistical matrix would be a positive identity matrix. However here the permutation matrix $F_{[1234]}$ and $F_{[1342]}$ above are obviously non-trivial matrix comparing with identity matrix. In fact, these permutation matrix form a group. when the spatial order of the particles returns to its initial ordering after a series of permutations, the product of those permutation matrices along the path should be equal to identity.

For a more general theoretical framework for permutating particles by Jordan-Wigner transformation, when a fermion moves one step forward, it would be multiplied by σ^z. If a fermion moves one step backward, a σ^z would be taken away from its string operator. To explicitly formulate this operation, we build ψ_μ fermion and b_μ fermion into a two dimensional vector,

$$\vec{f}_{\mathbf{i}_\mu} = (\psi_{\mathbf{i}_\mu}, b_{\mathbf{i}_\mu})^T. \tag{53}$$

There are two independent ways to switch the nearest neighboring two vectors of fermions operators. The first way is to map $\psi_{\mathbf{i}_{\mu-1}}$ to $\psi_{\mathbf{i}_\mu}$ and $b_{\mathbf{i}_{\mu-1}}$ to $b_{\mathbf{i}_\mu}$, the permutation matrix is

$$P_{\mathbf{i}_\mu,\mathbf{i}_{\mu-1}} = \begin{bmatrix} \sigma^z_{\mathbf{i}_{\mu-1}} & 0 \\ 0 & \sigma^z_{\mathbf{i}_{\mu-1}} \end{bmatrix}. \tag{54}$$

The other way is transforming $\psi_{\mathbf{i}_{\mu-1}}$ to $b_{\mathbf{i}_\mu}$ and mapping $b_{\mathbf{i}_{\mu-1}}$ to $\psi_{\mathbf{i}_\mu}$, the permutation matrix is off-diagonal matrix

$$P_{\mathbf{i}_\mu,\mathbf{i}_{\mu-1}} = \begin{bmatrix} 0 & -\sigma^z_{\mathbf{i}_{\mu-1}} \\ -\sigma^z_{\mathbf{i}_{\mu-1}} & 0 \end{bmatrix}. \tag{55}$$

Both the two matrices satisfy the algebra relation,

$$\vec{f}_{\mathbf{i}_\mu} = P_{\mathbf{i}_\mu,\mathbf{i}_{\mu-1}} \vec{f}_{\mathbf{i}_{\mu-1}}. \tag{56}$$

Every lattice site shoulders a pair of dual Majorana fermions. This is to compensate for the degree of freedom of spin operators. The example above suggest that only when we put all of the eight Majorana fermions into action, the Hilbert space generated by the fermion operators form a closed space. This suggest the eight Majorana fermion are highly entangled set of operators. This is another way to understand quantum entanglement.

VI. SUMMARY

The quantum statistics of exotic quasiparticles is more complicate than fermion and boson. It is a challenge to observe exotic quasiparticles in quantum optical experiment nowadays. We proposed a four particle model to study the explicit spin configuration of exotic quasiparticles which maybe is useful for quantum optical research.

This triangular star model is inspired by Kitaev honeycomb model. We introduced the spin coupling rules in Kitaev honeycomb model but add frustrated coupling terms between plaquette quasiparticles. Direct diagonalization of the Hamiltonian matrix produces the complete list of eigenenergy levels and eigenstates with explicit spin configuration. The eigenstates of this model are genuine entangled four qubit states. For instance, the first excited state can be viewed as a generalization of Greenberger-Horne-Zeilinger state and Yeo-Chua's genuine entangled states.

The quantum entanglement and quantum statistics for a clusters made of several-spins depends on how we pack the spins. We first take double-spin as a cluster. Usually a composite particle made of two spin 1/2 behaves like boson. This is indeed the case in zero energy level and the highest energy level where exchanging two double-spin clusters keeps the collective wave function invariant. However when we perform the same exchanging operation on ground state and intermediate excited state, there appears non-trivial off-diagonal elements in the statistical matrix. A double-spin 1/2 cluster may be a boson, a fermion, and other exotic particle. The zero energy state may be viewed as a singlet state of two double-spins. The highest excited state is a symmetric state of two double-spins.

The triple-spin cluster happened to be eigen-quasiparticles of this four spins model. We computed the statistical matrix for exchanging two triple-spin clusters on the first excited state, it does behave like a fermion as what we usually believe. The statistical matrix is a complicate matrix. So these triple-spin quasiparticles are neither boson nor fermions, they are exotic quasiparticles. Even in the zero energy state and the highest excited state where double-spin are normal bosons, the triple-spin cluster is still highly nontrivial, the von Neumann entropy of the reduced density operator of the triangular plaquette excitations is much higher than that of double-spin clusters. So the quantum entanglement in a triple-spin cluster is much stronger than that in a double spin cluster.



FIG. 4: . The extension of frustrated plaquette excitation model to Kitaev honeycomb model.

Permutation operation is the elementary operation for constructing quantum logical gate. Permutating Majorana fermions is believed to be a promising approach to quantum computation. However a free Majorana fermion is absent in nature. So we construct a composite Majorana fermion by a string of spins in the inverse Jordan-Wigner transformation. It fulfils the commutator relationship of Majorana fermions. The string operator of spins for Majorana fermion operator is different for different spatial ordering. So exchanging two Majorana fermions is encoded in the spatial order of spin representation. When we exchange two Majorana fermions, the effective statistical matrix is not simply adding a minus sign in front of the original wave funtion. It indicates the output states of a composite Majorana fermion is strongly entangled with other composite Majorana fermions. This statistical matrix is actually equivalent to permutation matrix. When the spatial order restored its original order after a circle of permutations, the product of these permutation matrices would be equal to identity. This operation constructs a permutation group.

The frustrated coupling term is the source of those exotic statistical behaviors. It can be straightly extended to Kitaev honeycomb model, we need to add antiferromagnetic coupling between the nearest neighboring plaquette upon the Kitaev honeycomb Hamiltonian H_k, $H = H_k + H_1$,

$$\begin{aligned} H_k &= \sum_{\mathbf{r}} (J_x \sigma^x_{\mathbf{r}} \sigma^x_{\mathbf{r}+\mathbf{e}_x} + J_y \sigma^y_{\mathbf{r}} \sigma^y_{\mathbf{r}+\mathbf{e}_y} + J_z \sigma^z_{\mathbf{r}} \sigma^z_{\mathbf{r}+\mathbf{e}_z}), \\ H_1 &= \sum_{<\mathbf{i}_p \mathbf{j}_p>} J_p S_{\mathbf{i}_p} S_{\mathbf{j}_p}, \quad S_{\mathbf{i}_p} = \sigma^z_1 \sigma^x_2 \sigma^y_3 \sigma^z_4 \sigma^x_5 \sigma^y_6, \end{aligned} \tag{57}$$

where $\mathbf{i}_p$ is the center of every plaquette(Fig. 4). All plaquette operators commute with Hamiltonian as well as themselves, $[S_{\mathbf{i}_p}, H] = 0$, $[S_{\mathbf{i}_p}, S_{\mathbf{j}_p}] = 0$. The newly added quasiparticle terms commute with Kitaev honeycomb Hamiltonian, $[H_k, H_1] = 0$. The plaquette coupling interaction is antiferromagnetic, i.e., $J_p > 0$. This model can be solved by inverse Jordan-Wigner transformation. A string operator of many spin operator is convenient for manipulating plaquette state. Further research will be carried out in the future.

VII. ACKNOWLEDGMENT

The author thanks Institute of Theoretical Physics-Chinese Academy of Sciences and Beijing Computation Science Research Center for hosting my visit.

VIII. APPENDIX

A. The inverse Jordan-Wigner transformation for different spatial ordering

The inverse Jordan-Wigner transformation for the ordering [1243] is

$$\begin{aligned} \psi_1 &= \sigma_1^y, \quad b_3 = -\sigma_3^y\sigma_1^z\sigma_4^z\sigma_2^z, \\ \psi_2 &= \sigma_2^y\sigma_1^z, \quad b_4 = -\sigma_4^y\sigma_1^z\sigma_2^z, \\ \psi_4 &= \sigma_4^x\sigma_1^z\sigma_2^z, \quad b_2 = -\sigma_2^x\sigma_1^z, \\ \psi_3 &= \sigma_3^x\sigma_1^z\sigma_4^z\sigma_2^z, \quad b_1 = -\sigma_1^x. \end{aligned} \tag{58}$$

The output state of the eight fermions for [1243] on ground state $|g\rangle_4$,

$$\begin{aligned} \psi_1|g\rangle_4 &= \frac{-i}{2}(|o\rangle_3 + |e\rangle_{10}), \ b_1|g\rangle_4 = \frac{1}{2}(|e\rangle_{12} - |g\rangle_1), \\ \psi_2|g\rangle_4 &= \frac{-i}{2}(|g\rangle_1 + |e\rangle_{12}), \ b_2|g\rangle_4 = \frac{1}{2}(|o\rangle_2 + |e\rangle_9), \\ \psi_4|g\rangle_4 &= -\frac{1}{2}(|g\rangle_3 + |e\rangle_{13}), \ b_4|g\rangle_4 = \frac{i}{2}(|o\rangle_2 - |e\rangle_9), \\ \psi_3|g\rangle_4 &= \frac{1}{2}(|o\rangle_3 - |e\rangle_{10}), \quad b_3|g\rangle_4 = \frac{i}{2}(|e\rangle_{13} - |g\rangle_3). \end{aligned} \tag{59}$$

Comparing the output states above with the output state for the initial spatial ordering [1423], we derived the permutation matrix for the spatial ordering [1243],

$$F_{[1243]} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -i & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & i & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -i & 0 & 0 & 0 & 0 \end{pmatrix}. \tag{60}$$

The inverse Jordan-Wigner transformation for the ordering [1234] is

$$\begin{aligned} \psi_1 &= \sigma_1^y, \quad b_4 = -\sigma_4^y\sigma_1^z\sigma_3^z\sigma_2^z, \\ \psi_2 &= \sigma_2^y\sigma_1^z, \quad b_3 = -\sigma_3^y\sigma_1^z\sigma_2^z, \\ \psi_3 &= \sigma_3^x\sigma_1^z\sigma_2^z, \quad b_2 = -\sigma_2^x\sigma_1^z, \\ \psi_4 &= \sigma_4^x\sigma_1^z\sigma_3^z\sigma_2^z, \quad b_1 = -\sigma_1^x. \end{aligned} \tag{61}$$

The output state of the eight fermions for [1234] on ground state $|g\rangle_4$,

$$\begin{aligned} \psi_1|g\rangle_4 &= \frac{-i}{2}(|o\rangle_3 + |e\rangle_{10}), \ \psi_2|g\rangle_4 = \frac{-i}{2}(|g\rangle_1 + |e\rangle_{12}), \\ \psi_3|g\rangle_4 &= \frac{-1}{2}(|g\rangle_3 - |e\rangle_{13}), \ \psi_4|g\rangle_4 = \frac{1}{2}(|o\rangle_2 - |e\rangle_9), \\ b_1|g\rangle_4 &= \frac{1}{2}(|e\rangle_{12} - |g\rangle_1), \quad b_2|g\rangle_4 = \frac{1}{2}(|o\rangle_2 + |e\rangle_9), \\ b_3|g\rangle_4 &= i\frac{1}{2}(|o\rangle_3 - |e\rangle_{10}), \quad b_4|g\rangle_4 = i\frac{1}{2}(|g\rangle_1 + |e\rangle_{12}). \end{aligned} \tag{62}$$

The inverse Jordan-Wigner transformation for the ordering [1324] is

$$\begin{aligned} \psi_1 &= \sigma_1^y, \quad b_4 = -\sigma_4^y\sigma_1^z\sigma_3^z\sigma_2^z, \\ \psi_3 &= \sigma_3^x\sigma_1^z, \quad b_2 = -\sigma_2^x\sigma_1^z\sigma_3^z, \\ \psi_2 &= \sigma_2^y\sigma_1^z\sigma_3^z, \quad b_3 = -\sigma_3^y\sigma_1^z, \\ \psi_4 &= \sigma_4^x\sigma_1^z\sigma_3^z\sigma_2^z, \quad b_1 = -\sigma_1^x. \end{aligned} \tag{63}$$

The output state of the eight fermions for [1324] on ground state $|g\rangle_4$,

$$\begin{aligned} \psi_1|g\rangle_4 &= \frac{-i}{2}(|o\rangle_3 + |e\rangle_{10}), \ \psi_3|g\rangle_4 = \frac{1}{2}(|o\rangle_3 - |e\rangle_{10}), \\ \psi_2|g\rangle_4 &= \frac{i}{2}(|o\rangle_2 + |e\rangle_9), \quad \psi_4|g\rangle_4 = \frac{1}{2}(|o\rangle_2 - |e\rangle_9), \\ b_1|g\rangle_4 &= \frac{1}{2}(|e\rangle_{12} - |g\rangle_1), \quad b_3|g\rangle_4 = \frac{i}{2}(|e\rangle_{13} - |g\rangle_3), \\ b_2|g\rangle_4 &= \frac{-1}{2}(|g\rangle_1 + |e\rangle_{12}), \ b_4|g\rangle_4 = \frac{-i}{2}(|e\rangle_{13} + |g\rangle_3). \end{aligned} \tag{64}$$

The inverse Jordan-Wigner transformation for the ordering [1342] is

$$\begin{aligned} \psi_1 &= \sigma_1^y, \quad b_2 = -\sigma_2^y\sigma_1^z\sigma_3^z\sigma_2^z, \\ \psi_3 &= \sigma_3^x\sigma_1^z, \quad b_4 = -\sigma_2^x\sigma_1^z\sigma_3^z, \\ \psi_4 &= \sigma_4^y\sigma_1^z\sigma_3^z, \quad b_3 = -\sigma_3^y\sigma_1^z, \\ \psi_2 &= \sigma_2^x\sigma_1^z\sigma_3^z\sigma_2^z, \quad b_1 = -\sigma_1^x. \end{aligned} \tag{65}$$

The output state of the eight fermions for [1342]on ground state $|g\rangle_4$,

$$\begin{aligned} \psi_1|g\rangle_4 &= \frac{-i}{2}(|o\rangle_3 + |e\rangle_{10}), \ \psi_3|g\rangle_4 = \frac{-1}{2}(|o\rangle_3 - |e\rangle_{10}), \\ \psi_4|g\rangle_4 &= \frac{i}{2}(|o\rangle_2 - |e\rangle_9), \quad \psi_2|g\rangle_4 = \frac{1}{2}(|o\rangle_2 + |e\rangle_9), \\ b_1|g\rangle_4 &= \frac{1}{2}(|e\rangle_{12} - |g\rangle_1), \quad b_3|g\rangle_4 = \frac{-i}{2}(|e\rangle_{13} - |g\rangle_3), \\ b_2|g\rangle_4 &= \frac{-i}{2}(|g\rangle_1 + |e\rangle_{12}), \ b_4|g\rangle_4 = \frac{-1}{2}(|e\rangle_{13} + |g\rangle_3). \end{aligned} \tag{66}$$

B. The 16 basis of spin configuration

The 16 basis for expanding the eigenvector of the 16 dimensional Hamiltonian matrix,

$$\begin{aligned}|1\rangle &= |\uparrow\uparrow\uparrow\uparrow\rangle, \; |2\rangle = |\uparrow\uparrow\uparrow\downarrow\rangle, \quad |3\rangle = |\uparrow\uparrow\downarrow\uparrow\rangle,\\ |4\rangle &= |\uparrow\uparrow\downarrow\downarrow\rangle, \; |5\rangle = |\uparrow\downarrow\uparrow\uparrow\rangle, \quad |6\rangle = |\uparrow\downarrow\uparrow\downarrow\rangle,\\ |7\rangle &= |\uparrow\downarrow\downarrow\uparrow\rangle, \; |8\rangle = |\uparrow\downarrow\downarrow\downarrow\rangle, \quad |9\rangle = |\downarrow\uparrow\uparrow\uparrow\rangle,\\ |10\rangle &= |\downarrow\uparrow\uparrow\downarrow\rangle, \; |11\rangle = |\downarrow\uparrow\downarrow\uparrow\rangle, \; |12\rangle = |\downarrow\uparrow\downarrow\downarrow\rangle,\\ |13\rangle &= |\downarrow\downarrow\uparrow\uparrow\rangle, \; |14\rangle = |\downarrow\downarrow\uparrow\downarrow\rangle, \; |15\rangle = |\downarrow\downarrow\downarrow\uparrow\rangle,\\ |16\rangle &= |\downarrow\downarrow\downarrow\downarrow\rangle. \qquad (67)\end{aligned}$$

[1] A. Kitaev, Ann. Phys. **321**, 2 (2006).

[2] C. Nayak, S. H. Simon, A. Stern, M. Freedman and S. Das Sarma, Rev. Mod. Phys. **80**, 1083 (2008).

[3] D. A. Ivanov, Phys. Rev. Lett. **86**, 268, (2001).

[4] X. Y. Feng, G. M. Zhang, and T. Xiang, Phys. Rev. Lett. **98**, 087204 (2007); H. D. Chen and J. P. Hu, Phys. Rev. B **76**, 193101 (2007).

[5] S. Dusuel, K. P. Schmidt, J. Vidal, Phys. Rev. Lett. **100**, 177204 (2008); G. Baskaran, S. Mandal and R. Shankar, Phys. Rev. Lett. **98**, 247201 (2007) .

[6] A. J. Willans, J. T. Chalker, and R. Moessner, Phys. Rev. Lett. **104**, 237203 (2010).

[7] Y. Yu and T. Si, arXiv:0804.0483 (2008).

[8] T. Si and Y. Yu, Nucl. Phys. B, **803**, 428 (2008).

[9] H. Yao, S. A. Kivelson, Phys. Rev. Lett. **99**, 247203 (2007).

[10] S. Yang, D. L. Zhou, and C. P. Sun, Phys. Rev. B **76**, 180404 (2007).

[11] Z. Wang, T. Ma, S. J. Gu, H. Q. Lin, Phys. Rev. A **81**, 062350 (2010).

[12] G. Kells, J. Kailasvuori, J. K. Slingerland, J. Vala, New J. Phys. **13**, 095014 (2011).

[13] Y. J. Han, R. Raussendorf, and L. M. Duan, Phys. Rev. Lett. **98**, 150404 (2007).

[14] C. Y. Lu, W. B. Gao,O. Guhne, X. Q. Zhou, Z. B. Chen, and J. W. Pan, Phys.Rev.Lett. **102**, 030502 (2009).

[15] C. Zhang, V. W. Scarola, S. Tewari and S. Das Sarma, Proc. Natl. Acad. Sci. USA **104**, 18415 (2007).

[16] M. Aguado, G. K. Brennen, F. Verstraete, and J. I. Cirac, Phys. Rev. Lett. **101**, 260501, (2008).

[17] H. Weimer, M. Muller, I. Lesanovsky, P. Zoller, and H. P. Buchler, Nature Phys. **6**, 382-388 (2010).

[18] H. Bombin, Phys. Rev. Lett. **105**, 030403, (2010); H. Bombin and M. A. Martin-Delgado, Phys. Rev. Lett. **98**, 160502 (2007).

[19] J. Chaloupka, G. Jackeli and G. Khaliullin, Phys. Rev. Lett. **105**, 027204 (2010); S. K. Choi, R. Coldea, A. N. Kolmogorov, T. Lancaster, I. I. Mazin, S. J. Blundell, P. G. Radaelli, Yogesh Singh, P. Gegenwart, K. R. Choi, S. W. Cheong, P. J. Baker, C. Stock, J. Taylor, Phys. Rev. Lett. **108**, 127204 (2012).

[20] C. W. J. Beenakker, Ann. Rev. Cond. Matt. Phys., arXiv:1112.1950, (2012).

[21] D. J. Clarke, J. Alicea, and K. Shtengel, arXiv: 1204.5479; J. Alicea, arXiv: 1202.1293., (2012).

[22] J. D. Sau, D. J. Clarke, and S. Tewari, Phys. Rev. B, **84**, 094505 (2011).

[23] K. Flensberg, Phys. Rev. Lett. **106**, 090503 (2011).

[24] A. Mezzacapo, J. Casanova, L. Lamata, E. Solano, arXiv:1111.5603, (2011).

[25] X. J. Liu, L. Jiang, H. Pu, and H. Hu, Phys. Rev. A, **85**, 021603(R) (2012).

[26] D. M. Greenberger, M.A. Horne, and A. Zeilinger, in Bells Theorem, Quantum Theory, and Conceptions of the Universe, edited by M. Kafatos (Kluwer Academic, Dordrecht, 1989), pp. 69C72.

[27] A. Zeilinger, M.A. Horne, and D. M. Greenberger, in Proceedings of Squeezed States and Quantum Uncertainty, edited by D. Han, Y. S. Kim, and W.W. Zachary, NASA Conference Publication No. 3135 (NASA, Washington, DC, 1992), pp. 73C81.

[28] Y. Yeo and W. K. Chua, Phys. Rev. Lett. **84**, 4457 (2000).

[29] L. F. Zhao , B. H. Lai , F. Mei Y. F. Yu, X. L. Feng and Z. M. Zhang, Chin. Phys. B **19**, 094207 (2010).

[30] R. Moessner, S. L. Sondhi and P. Chandra, Phys. Rev. Lett. **84**, 4457 (2000); R. Moessner and S. L. Sondhi, arXiv:cond-mat/0011250, (2000).

[31] M. Y. Ye and X. M. Lin, Phys. Let. A **372** 4157, (2008).

[32] W. K. Wootters, Phys. Rev. Lett. 80, 2245, (1998); Hill S. and Wootters W. K., Phys. Rev. Lett. 78 5022, (1997).

[33] C. D. Batista, G. Ortiz, Phys. Rev. Lett. **86**, 1082 (2001); M. S. Kochmanski, J. Tech. Phys., 36, 485 (1995), arXiv:cond-mat/9807388.